

Minimal Length and Generalized Dirac Equation

Kourosh Nozari and **Mojdeh Karami**

*Department of Physics, Faculty of Basic Science,
University of Mazandaran,
P. O. Box 47416-1467, Babolsar, IRAN
e-mail: knozari@umz.ac.ir*

Abstract

Existence of a minimal observable length which has been indicated by string theory and quantum gravity, leads to a modification of Dirac equation. In this letter we find this modified Dirac equation and solve its eigenvalue problem for a free particle. We will show that due to background spacetime fluctuation, it is impossible to have free particle in Planck scale.

PACS: 03.65.Pm, 04.60.-m

Keywords: Quantum Gravity, Generalized Uncertainty Principle, Relativistic Quantum Mechanics, Dirac Equation

1 Introduction

The problem of reconciling Quantum Mechanics with General Relativity is one of the task of modern theoretical physics which, until now, has not yet found a consistent and satisfactory solution. One of the most interesting consequences of this unification is that in quantum gravity there exists a minimal observable distance on the order of the Planck distance, $l_P = \sqrt{\frac{G\hbar}{c^3}} \sim 10^{-33}cm$, where G is the Newton constant. The existence of such a fundamental length is a dynamical phenomenon due to the fact that, at Planck scale, there are fluctuations of the background metric, i.e. a limit of the order of Planck length appears when quantum fluctuations of the gravitational field are taken into account. In the language of string theory one can say that a string cannot probe distances smaller than its length. The existence of such minimal observable length which is motivated from string theory[1-5], loop quantum gravity[6], Non-commutative geometry[7] and black hole physics[8,9], leads to a generalization of Heisenberg uncertainty principle to incorporate gravitational induced uncertainty from very beginning. This generalized uncertainty principle can be written as

$$\Delta x \geq \frac{\hbar}{\Delta p} + \alpha' l_P^2 \frac{\Delta p}{\hbar}. \quad (1)$$

At energy much below the Planck mass, $m_P = \sqrt{\frac{\hbar c}{G}} \sim 10^{19}GeV/c^2$, the extra term in equation (1) is irrelevant and the Heisenberg uncertainty relation is recovered, while, as we approach the Planck energy, this term becomes relevant and, as has been said, it is related to the existence of minimal observable length. Now the generalized commutation relation becomes,

$$[x, p] = i\hbar(1 + \beta' p^2), \quad (2)$$

where $\beta' = \alpha' \frac{l_P^2}{\hbar^2}$. This feature constitutes a part of the motivation to study the effects of this modified algebra on various observable. Consequences of such a gravitational uncertainty principle (GUP), has been studied extensively[10-25]. Here we proceed some more step in this direction. We consider the effect of GUP on definition of momentum operator and then we obtain a generalized Dirac equation and solve the corresponding eigenvalue problem. We will show that it is impossible to have free particle in Planck scale. Comparison between our results and the results of ordinary relativistic quantum mechanics shows a considerable departure when one approaches extreme quantum gravity limit. We choose $\hbar = c = 1$. In this situation $\beta' = l_P^2 = \frac{1}{m_P^2}$. Latin index i , takes the values 1, 2, 3 and Greek index μ takes the values 0, 1, 2, 3. Index 0 stands for temporal part of 4-vectors.

2 Generalized Dirac Equation

Using GUP as equation (1), one can generalize usual momentum operator $p_{op} = -i\frac{\partial}{\partial x}$ to the following form[19,20]

$$p_{op}^{(GUP)} = -i\left(1 - \beta'\left(\frac{\partial}{\partial x}\right)^2\right)\frac{\partial}{\partial x}, \quad (3)$$

or in three-dimensional form,

$$\vec{p} = -i\vec{\nabla}\left(1 - \beta'\vec{\nabla}^2\right). \quad (4)$$

To satisfy Lorentz invariance, this equation can be written as

$$p_i = -i\left[1 - \beta'(\partial_i)^2\right]\partial_i. \quad (5)$$

The square of this operator is,

$$p^2 = p_i p^i = \left(-i\left[1 - \beta'(\partial_i)^2\right]\partial_i\right)\left(-i\left[1 - \beta'(\partial^i)^2\right]\partial_i\right), \quad (6)$$

which up to first order in β' simplifies as

$$p^2 \simeq -\left[\partial_i \partial^i - 2\beta'(\partial^i \partial_i)(\partial^i \partial_i)\right]. \quad (7)$$

Now the Dirac equation[26],

$$(\gamma^\mu p_\mu - m)\psi = (i\gamma^\mu \partial_\mu + m)\psi = 0, \quad (8)$$

which can be written as

$$i\partial_0\psi = -\gamma_0(i\gamma^i \partial_i + m)\psi, \quad (9)$$

should be written in generalized form. For this purpose, in addition to generalization of momentum operator, we should consider the generalization of energy operator also. Usually one considers the energy operator as $E = i\partial_0$. As a consequence of generalized uncertainty principle, one can define generalized frequency as[27]

$$\tilde{\omega} = E - \beta'E^3 = E(1 - \beta'E^2), \quad (10)$$

where a numerical factor of $1/3$, which is irrelevant to our purposes has been omitted. Using the energy mass shell condition $p^2 + m^2 = E^2$, one finds

$$\tilde{E} = E\left(1 - \beta'(p^2 + m^2)\right) = E\left(1 - \beta'p^2 - \beta'm^2\right). \quad (11)$$

Now the generalized Dirac equation can be written as

$$i\partial_0\tilde{\psi} = -\gamma_0(i\gamma^i\partial_i + m)(1 - \beta'(p_ip^i) - \beta'm^2)\tilde{\psi}. \quad (12)$$

Since p_ip^i is given by equation (7), we find the following form for generalized Dirac equation

$$i\partial_0\tilde{\psi} = -\gamma_0(i\gamma^i\partial_i + m)\left[1 + \beta'(\partial_i\partial^i - m^2)\right]\tilde{\psi}, \quad (13)$$

up to first order in β' . In this equation $\tilde{\psi}$ is generalized Dirac spinor which will be calculated. Note that this equation has Lorentz invariance evidently. To solve this eigenvalue equation, first we try to write it in a form which be comparable easily with usual Dirac equation, (9). For this purpose, the right hand side of equation (13) can be written as,

$$\begin{aligned} & -\gamma_0(i\gamma^i\partial_i + m)(1 + \beta'\partial^i\partial_i - \beta'm^2)\tilde{\psi}, \\ & = -\gamma_0(i\gamma^i\partial_i + i\beta'\gamma^i(\partial^i\partial_i)\partial_i - \beta'm^2i\gamma^i\partial_i + m + \beta'm\partial_i\partial^i - \beta'm^3)\tilde{\psi}. \end{aligned} \quad (14)$$

Therefore, equation (13) can be re-written as

$$i\partial_0\tilde{\psi} = -\gamma_0(i\gamma^i\partial_i + i\beta'\gamma^i(\partial^i\partial_i)\partial_i - \beta'm^2i\gamma^i\partial_i + m + \beta'm\partial_i\partial^i - \beta'm^3)\tilde{\psi}, \quad (15)$$

which leads to

$$\left[E + [\beta\beta'm(p^2 - m^2)]\right]\tilde{\psi} = \left[(\alpha\cdot\mathbf{p})(1 - \beta'(\nabla^2 + m^2) + \beta m)\right]\tilde{\psi}. \quad (16)$$

In comparison with equation (9), here we define a "generalized potential"

$$V_0 = \beta'm(p^2 - m^2) \quad (17)$$

and "generalized momentum"

$$\tilde{\Pi} = p(1 - \beta'(\nabla^2 + m^2)). \quad (18)$$

Note that when $\beta' \rightarrow 0$, one finds $V_0 \rightarrow 0$ and this shows that V_0 can be interpreted as a potential related to quantum gravitational effects. In the presence of this generalized potential the particle is no longer free. Therefore in Planck scale there is no free particle due to quantum fluctuations of background spacetime. Now the generalized Hamiltonian becomes,

$$\tilde{H} = (\alpha\cdot\mathbf{p})(1 - \beta'(\nabla^2 + m^2)) + \beta m \quad (19)$$

and therefore generalized Dirac equation can be written as

$$(E + \beta V_0)\tilde{\psi} = \tilde{H}\tilde{\psi}. \quad (20)$$

To solve this generalized Dirac equation for "free particle", we consider the following solution

$$\tilde{\psi}(z, t) = \tilde{\psi}(z)e^{-i\varepsilon t}, \quad (21)$$

where $\varepsilon = \lambda(E + V_0)$ and we have assumed that the particle is moving in z -direction. $\lambda = +1$ or -1 is related to particle and anti-particle spectrum respectively. We consider the case $\lambda = +1$ in which follows. Now, regarding to the 4-dimensional nature of spinors, we have

$$\tilde{\psi}(z) = \begin{pmatrix} \varphi \\ \chi \end{pmatrix} = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \end{pmatrix}. \quad (22)$$

Using operator form of $\tilde{H}$ as equation (19), we find

$$\left[E \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + V_0 \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right] \begin{pmatrix} \varphi \\ \chi \end{pmatrix} = \begin{pmatrix} \mathbf{0} & \sigma \\ \sigma & \mathbf{0} \end{pmatrix} \cdot \mathbf{p} \left(1 - \beta'(\nabla^2 + m^2) \right) \begin{pmatrix} \varphi \\ \chi \end{pmatrix} + m \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \varphi \\ \chi \end{pmatrix}. \quad (23)$$

We should find eigenvalues and eigenvectors of this equation. Suppose that

$$\varphi(z) = \varphi_0 e^{ip(1-\beta'p^2)z} \quad (24)$$

and

$$\chi(z) = \chi_0 e^{ip(1-\beta'p^2)z}. \quad (25)$$

Two points should be notified here: first note that we have considered the generalized form of momentum operator in these equations. Secondly, existence of minimal length destroys the notion of locality. As Kempf has shown, in this situation one should consider maximally localized states[25]. Therefore (24) and (25) are maximally localized states. After substituting these two supposed solutions in equation (23), some simple algebra gives

$$\begin{aligned} (E + V_0)\varphi_0 &= (\sigma_z \cdot p)[1 - \beta'(2p^2 + m^2)]\chi_0 + m\varphi_0 \\ (E - V_0)\chi_0 &= (\sigma_z \cdot p)[1 - \beta'(2p^2 + m^2)]\varphi_0 - m\chi_0, \end{aligned} \quad (26)$$

where leads to

$$\chi_0 = \frac{(\sigma_z \cdot p)(1 - \beta'(2p^2 + m^2))}{E - V_0 + m} \varphi_0. \quad (27)$$

Now suppose that

$$\varphi_0 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad (28)$$

then it follows that

$$\chi_0 = \frac{(\sigma_z \cdot p) \left(1 - \beta' (2p^2 + m^2) \right)}{E - V_0 + m} \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \quad (29)$$

To find energy spectrum of the system one should evaluate the following 4×4 determinant

$$\begin{vmatrix} E + V_0 - m & -(\sigma_z \cdot p) \left(1 - \beta' (2p^2 + m^2) \right) \\ -(\sigma_z \cdot p) \left(1 - \beta' (2p^2 + m^2) \right) & E - V_0 + m \end{vmatrix} = 0. \quad (30)$$

Using the identity

$$(\sigma \cdot \mathbf{A})(\sigma \cdot \mathbf{B}) = \mathbf{A} \cdot \mathbf{B} + i\sigma \cdot (\mathbf{A} \times \mathbf{B}) \quad (31)$$

a simple calculation gives

$$E^2 = p^2 + m^2 + [2\beta'(2p^4 - p^2m^2 + m^4)] \quad (32)$$

up to first order in β' . The term in curly bracket gives the correction due to quantum gravitational effects. In usual quantum mechanics, $\beta' \ll 1$, and therefore one recovers the well-known result for relativistic free particle. In extreme quantum gravity limit when $\beta' \sim 1$, there is a considerable departure from usual picture.

Now the time-dependent wave function of the system becomes,

$$\tilde{\psi} = N e^{i[p(1-\beta'p^2)z - \lambda(E+V_0)t]} \left(\frac{\varphi}{\frac{(\sigma_z \cdot p) \left(1 - \beta' (2p^2 + m^2) \right)}{E + m - V_0} \varphi} \right), \quad (33)$$

or equivalently

$$\tilde{\psi} = N e^{i[p(1-\beta'p^2)z - \lambda(E+V_0)t]} \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix} \frac{(\sigma_z \cdot p) \left(1 - \beta' (2p^2 + m^2) \right)}{E + m - V_0} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right). \quad (34)$$

The normalization constant can be calculated using orthogonality condition

$$\int \tilde{\psi}_{\lambda,p}^\dagger(z, t) \tilde{\psi}_{\lambda,p'}(z, t) d^3z = \delta_{\lambda,\lambda'} \delta(p - p'), \quad (35)$$

which leads to

$$N^2 \left(\varphi^\dagger \varphi + \varphi^\dagger \left(\frac{(\sigma_z \cdot p) \left(1 - \beta' (2p^2 + m^2) \right)}{E + m - V_0} \right)^2 \varphi \right) = 1 \quad (36)$$

or

$$N = \sqrt{\frac{(E + m - V_0)^2}{(E + m - V_0)^2 + \left((\sigma_z \cdot p)(1 - \beta'(2p^2 + m^2))\right)^2}}. \quad (37)$$

In terms of explicit form of V_0 and $\tilde{\Pi}$, and using equation (31), this normalization factor becomes

$$N = \sqrt{\frac{p^2 + 2m^2 + 4\beta'p^4 + 2Em}{2m^2 + 2p^2 - 2\beta'p^2m^2 + 2Em}}. \quad (38)$$

Now the complete wave function for "free particle" in GUP is given by

$$\tilde{\psi}_{p,\lambda}(z, t) = \sqrt{\frac{p^2 + 2m^2 + 4\beta'p^4 + 2Em}{2m^2 + 2p^2 - 2\beta'p^2m^2 + 2Em}} \begin{pmatrix} 1 \\ 0 \\ \frac{(\sigma_z \cdot p)(1 - \beta'(2p^2 + m^2))}{E + m - V_0} \\ 0 \end{pmatrix} e^{i[p(1 - \beta'p^2)z - \lambda(E + V_0)t]} \quad (39)$$

for $\lambda = \pm 1$. In the limit of $\beta' \rightarrow 0$, this equation reduces to usual Dirac spinor for free particle[26]. Now one should consider Helicity. In relativistic quantum mechanics, Helicity is defined as[26]

$$\Lambda_s = \Sigma \cdot \hat{\mathbf{p}}, \quad (40)$$

where

$$\Sigma = \begin{pmatrix} \sigma & \mathbf{0} \\ \mathbf{0} & \sigma \end{pmatrix}, \quad (41)$$

and $\hat{\mathbf{p}} = \frac{\mathbf{p}}{p}$. When one considers the generalized form of momentum operator, the $\hat{\mathbf{p}}$ unit vector does not change and therefore, Helicity operator suffers no change in GUP. It is straightforward to show that generalized Hamiltonian and generalized momentum operator commute with Helicity operator, i.e. $[\tilde{H}, \Lambda_s] = 0$ and $[\tilde{\Pi}, \Lambda_s] = 0$. Now we should consider this further degree of freedom in complete wave function. Since we have considered the z -direction as our preferred direction, there are two possible eigenvalues for Helicity operator: $\pm 1/2$. Therefore the complete Dirac spinor in the framework of GUP for "free particle" is given by

$$\tilde{\psi}_{p,\lambda,+\frac{1}{2}} = \sqrt{\frac{p^2 + 2m^2 + 4\beta'p^4 + 2Em}{2m^2 + 2p^2 - 2\beta'p^2m^2 + 2Em}} \begin{pmatrix} 1 \\ 0 \\ \frac{(\sigma_z \cdot p)(1 - \beta'(2p^2 + m^2))}{E + m - V_0} \\ 0 \end{pmatrix} e^{i[p(1 - \beta'p^2)z - \lambda(E + V_0)t]}, \quad (42)$$

for positive Helicity and

$$\tilde{\psi}_{p,\lambda,-\frac{1}{2}} = \sqrt{\frac{p^2 + 2m^2 + 4\beta'p^4 + 2Em}{2m^2 + 2p^2 - 2\beta'p^2m^2 + 2Em}} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \left(\frac{(\sigma_i \cdot p)(1 - \beta'(2p^2 + m^2))}{E + m - V_0} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) e^{i[p(1 - \beta'p^2)z - \lambda(E + V_0)t]}, \quad (43)$$

for negative Helicity. The orthogonality condition,

$$\int \tilde{\psi}_{p_z,\lambda,s_z}^\dagger(z,t) \tilde{\psi}_{p'_z,\lambda,s'_z}(z,t) d^3x = \delta_{\lambda,\lambda'} \delta_{s_z,s'_z} \delta(p_z - p'_z) \quad (44)$$

is satisfied properly.

3 Summary

Gravity induces uncertainty. Considering this additional gravitational induced uncertainty, usual uncertainty principle of Heisenberg should be modified. This fact leads to generalized uncertainty principle. An immediate consequence of this generalized uncertainty principle is generalization of momentum operator. This generalized form of momentum operator should leads us to a generalized Dirac equation. In this paper we have found this generalized Dirac equation and have solved its eigenvalue problem. We have shown that in Planck scale, due to quantum fluctuation of background spacetime, there is no free particle. This have led us to define a quantum gravitational generalized potential. Our calculation shows that Helicity operator do not changes in GUP and therefore complete generalized spinor of "free particle" has been found systematically. Comparison between our results with usual result of relativistic particle shows that in extreme quantum gravity limit, $\beta' \rightarrow 0$, the departure from ordinary results are considerable.

References

- [1] G. Veneziano, Europhys. Lett. 2 (1986) 199
- [2] D. Amati, M. Ciafaloni, and G. Veneziano, Phys. Lett. B197 (1987) 81
- [3] K. Konishi, G. Paffuti, and P. Provero, Phys. Lett. B234 (1990) 276
- [4] M. Kato, Phys. Lett. B245 (1990) 43

- [5] R. Guida, K. Konishi, and P. Provero, Mod. Phys. Lett. A6 (1991) 1487
- [6] L. J. Garay, Int. J. Mod. Phys. A10 (1995) 145
- [7] S. Capozziello, G. Lambiase, G. Scarpetta, Int.J.Theor.Phys. 39 (2000) 15
- [8] F. Scardigli, Phys. Lett. B452 (1999) 39-44
- [9] F. Scardigli and R. Casadio, Class. Quant. Grav. 20 (2003) 3915-3926
- [10] M. Maggiore, Phys. Lett. B304 (1993) 65
- [11] C. Castro, Found. Phys. Lett. 10 (1997) 273.
- [12] A. Camacho, Gen. Rel. Grav. 34 (2002) 1839.
- [13] M. Maggiore, Phys. Rev. D49 (1994) 5182.
- [14] M. Maggiore, Phys. Lett. B319 (1993) 83.
- [15] S. Kalyana Rama, Phys.Lett. B 519 (2001) 103.
- [16] A. Camacho, Gen. Rel. Grav. 35 (2003) 1153.
- [17] F. Scardigli, R. Casadio, Class. Quantum Grav. 20 (2003) 3915
- [18] S. Hossenfelder, Mod.Phys.Lett. A19 (2004) 2727-2744.
S. Hossenfelder, Phys.Rev. D70 (2004) 105003
- [19] F. Brau, J.Phys. A32 (1999) 7691-7696
- [20] R. Akhoury and Y. P. Yao, Phys.Lett. B572 (2003) 37-42
- [21] R. J. Adler, P. Chen, D. I. Santiago, Gen. Rel. Grav. 33 (2001) 2101
- [22] R. Adler and D. Santiago, Mod. Phys. Lett. A14 (1999) 1371
- [23] R. Adler and D. Santiago, arXiv: hep-th/9908073
- [24] K. Nozari and S. M. Mehdipour, to be appear in Mod. Phys. Lett. A, 2005
- [25] A. Kempf *et al*, Phys. Rev. D52 (1995) 1108

- [26] S. Hossenfelder *et al*, Phys. Lett. B 575 (2003) 85-99
- [27] W. Greiner, *Relativistic Quantum Mechanics: Wave Equation*, Springer-Verlag, 2000